– Ask the patient: ————————————————————————————————————		
	OVes	ONa
1. In the past few weeks, have you wished you were dead?	O Yes	O No
2. In the past few weeks, have you felt that you or your family would be better off if you were dead?	<b>○</b> Yes	O No
3. In the past week, have you been having thoughts about killing yourself?	O Yes	O No
4. Have you ever tried to kill yourself?	O Yes	O No
If yes, how?		
When?		
If the patient answers Yes to any of the above, ask the following acuit	ty question:	
5. Are you having thoughts of killing yourself right now?	<b>O</b> Yes	ONo
If yes, please describe:		
Next steps:		
<ul> <li>If patient answers "No" to all questions 1 through 4, screening is complete (not necessary No intervention is necessary (*Note: Clinical judgment can always override a negative screen,</li> </ul>		
<ul> <li>If patient answers "Yes" to any of questions 1 through 4, or refuses to answer, they are c positive screen. Ask question #5 to assess acuity:</li> </ul>	onsidered a	
<ul> <li>"Yes" to question #5 = acute positive screen (imminent risk identified)</li> <li>Patient requires a STAT safety/full mental health evaluation.</li></ul>	ın or clinician	
<ul> <li>"No" to question #5 = non-acute positive screen (potential risk identified)</li> <li>Patient requires a brief suicide safety assessment to determine if a full ment is needed. Patient cannot leave until evaluated for safety.</li> <li>Alert physician or clinician responsible for patient's care.</li> </ul>	al health evaluation	,

## Provide resources to all patients -

- 24/7 National Suicide Prevention Lifeline 1-800-273-TALK (8255) En Español: 1-888-628-9454
- 24/7 Crisis Text Line: Text "HOME" to 741-741